

Twistor spinors on Lorentzian symmetric spaces

Helga Baum

February 1, 2008

Abstract

We solve the twistor equation on all indecomposable Lorentzian symmetric spaces explicitly.

1 Introduction

Let (M^n, g) be an oriented semi-Riemannian spin manifold with the spinor bundle S . The *twistor operator* $\mathcal{D}$ is defined as the composition of the spinor derivative ∇^S with the projection p onto the kernel of the Clifford multiplication μ

$$\mathcal{D} : \Gamma(S) \xrightarrow{\nabla^S} \Gamma(T^*M \otimes S) \xrightarrow{g} \Gamma(TM \otimes S) \xrightarrow{p} \Gamma(\ker \mu).$$

The solutions of the conformally invariant equation $\mathcal{D}\varphi = 0$ are called *twistor spinors*. Twistor spinors were introduced by R.Penrose in General Relativity (see [Pen67], [PR86], [NW84]). They are related to Killing vector fields in semi-Riemannian supergeometry (see [ACDS97]). In the last years essential results concerning the geometry of Riemannian spin manifolds admitting twistor spinors were obtained by A.Lichnerowicz, T.Friedrich, K.Habermann, H-B.Rademacher, W.Kuehnel and other authors. For a survey on the literature cf. [Fri97]. In the Lorentzian setting there was established a relation between a special class of solutions of the twistor equation and the Fefferman spaces occurring in CR-geometry (cf. [Lew91], [Bau97]).

Let $\mathcal{T}(M^n, g)$ denote the space of all twistor spinors of (M, g) . It is known that

$$\dim \mathcal{T}(M^n, g) \leq 2 \cdot 2^{\lfloor \frac{n}{2} \rfloor}$$

(see [BFGK91]). If (M^n, g) is conformally flat and simply connected, then one has $\dim \mathcal{T}(M^n, g) = 2 \cdot 2^{\lfloor \frac{n}{2} \rfloor}$. In the present paper we determine the twistor spinors on all indecomposable Lorentzian symmetric spaces explicitly. In particular, we prove:

1. If (M^n, g) is an indecomposable non-conformally flat Lorentzian symmetric spin manifold of dimension $n \geq 3$, then each twistor spinor is parallel and $\dim \mathcal{T}(M^n, g) = q \cdot 2^{\lfloor \frac{n}{2} \rfloor}$, where $q = \frac{1}{2}, \frac{1}{4}$ or 0, depending on the fundamental group $\pi_1(M)$ and on the spin structure of (M^n, g) .
2. If (M^n, g) is an indecomposable conformally flat Lorentzian symmetric spin manifold of dimension $n \geq 3$ and non-constant sectional curvature, then $\dim \mathcal{T}(M^n, g) = q \cdot 2^{\lfloor \frac{n}{2} \rfloor}$, where $q = 2, \frac{3}{2}, 1, \frac{3}{4}$ or 0, depending on $\pi_1(M)$ and on the spin structure.

3. If (M^n, g) is a Lorentzian symmetric spin manifold of dimension $n \geq 3$ and constant sectional curvature, then $\dim \mathcal{T}(M^n, g) = q \cdot 2^{\lfloor \frac{n}{2} \rfloor}$, where $q = 2, 1$, or 0 , depending on $\pi_1(M)$ and on the spin structure.

The calculations are based on the fact, that the indecomposable Lorentzian symmetric spaces are completely classified (see [CW70]).

2 Lorentzian symmetric spaces

Let us first recall the description of Lorentzian symmetric spaces. A connected semi-Riemannian manifold (M, g) is called *indecomposable*, if there is no proper, nondegenerate subspace of $T_x M$ invariant under the action of the holonomy group $\text{Hol}_x(g)$. Each simply connected semi-Riemannian symmetric space is isometric to a product $M_0 \times M_1 \times \dots \times M_r$, where M_i , $i = 1, \dots, r$, are indecomposable simply connected semi-Riemannian symmetric spaces of dimension ≥ 2 and M_0 is semi-Euclidean.

Let (M^n, g) be a Lorentzian symmetric space. By $G(M)$ we denote the group of transvections of (M^n, g) and by $\mathfrak{g}$ its Lie algebra. One has the following structure result:

Theorem 1 ([CW70])

Let (M^n, g) be an indecomposable Lorentzian symmetric space of dimension $n \geq 2$. Then the Lie algebra $\mathfrak{g}$ of the transvection group of (M^n, g) is either semi-simple or solvable.

Let $\underline{\lambda} = (\lambda_1, \dots, \lambda_{n-2})$ be an $(n-2)$ -tupel of real numbers $\lambda_j \in \mathbb{R} \setminus \{0\}$ and let us denote by $M_{\underline{\lambda}}^n$ the Lorentzian space $M_{\underline{\lambda}}^n := (\mathbb{R}^n, g_{\underline{\lambda}})$, where

$$(g_{\underline{\lambda}})_{(s,t,x)} := 2ds dt + \sum_{j=1}^{n-2} \lambda_j x_j^2 ds^2 + \sum_{j=1}^{n-2} dx_j^2.$$

If $\underline{\lambda}_\pi = (\lambda_{\pi(1)}, \dots, \lambda_{\pi(n-2)})$ is a permutation of $\underline{\lambda}$ and $c > 0$, then $M_{\underline{\lambda}}^n$ is isometric to $M_{c\underline{\lambda}_\pi}^n$.

Theorem 2 ([CW70], [CP80])

Let (M^n, g) be an indecomposable solvable Lorentzian symmetric space of dimension $n \geq 3$. Then (M^n, g) is isometric to $M_{\underline{\lambda}}^n/A$, where $\underline{\lambda} \in (\mathbb{R} \setminus \{0\})^{n-2}$ and A is a discrete subgroup of the centralizer $Z_{I(M_{\underline{\lambda}})}(\bar{G}(M_{\underline{\lambda}}))$ of the transvection group $G(M_{\underline{\lambda}})$ in the isometry group $I(M_{\underline{\lambda}})$ of $M_{\underline{\lambda}}^n$.

For the centralizer $Z_{\underline{\lambda}} := Z_{I(M_{\underline{\lambda}})}(G(M_{\underline{\lambda}}))$ it is known:

Theorem 3 ([CK78])

Let $\underline{\lambda} = (\lambda_1, \dots, \lambda_{n-2})$ be a tupel of non-zero real numbers.

1. *If there is a positive λ_i or if there are two numbers λ_i, λ_j such that $\frac{\lambda_i}{\lambda_j} \notin \mathbb{Q}^2$, then $Z_{\underline{\lambda}} \simeq \mathbb{R}$ and $\varphi \in Z_{\underline{\lambda}}$ if and only if $\varphi(s, t, x) = (s, t + \alpha, x)$, $\alpha \in \mathbb{R}$.*

2. Let $\lambda_i = -k_i^2 < 0$ and $\frac{k_i}{k_j} \in \mathbb{Q}$ for all $i, j \in \{1, \dots, n-2\}$. Then $\varphi \in Z_\Delta$ if and only if

$$\varphi(s, t, x) = (s + \beta, t + \alpha, (-1)^{m_1} x_1, \dots, (-1)^{m_{n-2}} x_{n-2}),$$

where $\alpha \in \mathbb{R}$, $m_1, \dots, m_{n-2} \in \mathbb{Z}$ and $\beta = \frac{m_i \cdot \pi}{k_i}$ for all $i = 1, \dots, n-2$.

Let us denote by $S_1^n(r)$ the pseudo-sphere

$$S_1^n(r) := \{x \in \mathbb{R}^{n+1,1} \mid \langle x, x \rangle_{n+1,1} = -x_1^2 + x_2^2 + \dots + x_{n+1}^2 = r^2\} \subset \mathbb{R}^{n+1,1}$$

and by $H_1^n(r)$ the pseudo-hyperbolic space

$$H_1^n(r) := \{x \in \mathbb{R}^{n+1,2} \mid \langle x, x \rangle_{n+1,2} = -x_1^2 - x_2^2 + x_3^2 + \dots + x_{n+1}^2 = -r^2\} \subset \mathbb{R}^{n+1,2}$$

with the Lorentzian metrics induced by $\langle \cdot, \cdot \rangle_{n+1,1}$ and $\langle \cdot, \cdot \rangle_{n+1,2}$, respectively.

Theorem 4 ([CLP⁺90], [Wol84])

Let (M^n, g) be an indecomposable semi-simple Lorentzian symmetric space of dimension $n \geq 3$. Then (M^n, g) has constant sectional curvature $k \neq 0$. Therefore, it is isometric to $S_1^n(r)/\{\pm I\}$ or $S_1^n(r)$, ($k = \frac{1}{r^2} > 0$), or to a Lorentzian covering of $H_1^n(r)/\{\pm I\}$, ($k = -\frac{1}{r^2} < 0$).

3 Spinor representation

For concrete calculations we will use the following realization of the spinor representation. Let $\text{Cliff}_{n,k}$ be the Clifford algebra of $(\mathbb{R}^n, -\langle \cdot, \cdot \rangle_k)$, where $\langle \cdot, \cdot \rangle_k$ is the scalar product $\langle x, y \rangle_k := -x_1 y_1 - \dots - x_k y_k + x_{k+1} y_{k+1} + \dots + x_n y_n$. For the canonical basis $(e_1, \dots, e_n)$ of $\mathbb{R}^n$ one has the following relations in $\text{Cliff}_{n,k}$: $e_i \cdot e_j + e_j \cdot e_i = -2\varepsilon_j \delta_{ij}$,

where $\varepsilon_j = \begin{cases} -1 & j \leq k \\ 1 & j > k \end{cases}$. Denote $\tau_j = \begin{cases} i & j \leq k \\ 1 & j > k \end{cases}$ and

$$U = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad V = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad E = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad T = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}.$$

If $n = 2m$ is even, we have an isomorphism

$$\phi_{2m,k} : \text{Cliff}_{2m,k}^{\mathbb{C}} \xrightarrow{\sim} M(2^m; \mathbb{C})$$

given by the Kronecker product

$$\begin{aligned} \phi_{2m,k}(e_{2j-1}) &= \tau_{2j-1} & E \otimes \dots \otimes E \otimes U \otimes T \otimes \dots \otimes T \\ \phi_{2m,k}(e_{2j}) &= \tau_{2j} & E \otimes \dots \otimes E \otimes V \otimes \underbrace{T \otimes \dots \otimes T}_{j-1}. \end{aligned} \quad (1)$$

If $n = 2m + 1$ is odd and $k < n$, we have the isomorphism

$$\phi_{2m+1,k} : \text{Cliff}_{2m+1,k}^{\mathbb{C}} \xrightarrow{\sim} M(2^m; \mathbb{C}) \oplus M(2^m; \mathbb{C})$$

given by

$$\begin{aligned}\Phi_{2m+1,k}(e_j) &= (\Phi_{2m,k}(e_j), \Phi_{2m,k}(e_j)), \quad j = 1, \dots, 2m \\ \Phi_{2m+1,k}(e_n) &= \tau_n(i T \otimes \dots \otimes T, -i T \otimes \dots \otimes T).\end{aligned}\tag{2}$$

Let $\text{Spin}(n, k) \subset \text{Cliff}_{n,k}$ be the spin group. The spinor representation is given by

$$\kappa_{n,k} = \hat{\phi}_{n,k}|_{\text{Spin}(n,k)} : \text{Spin}(n, k) \longrightarrow \text{GL}(\mathbb{C}^{2^{\lfloor \frac{n}{2} \rfloor}}),$$

where $\hat{\phi}_{2m,k} = \phi_{2m,k}$ and $\hat{\phi}_{2m+1,k} = pr_1 \circ \phi_{2m+1,k}$. We denote this representation by $\Delta_{n,k}$. If $n = 2m$, $\Delta_{2m,k}$ splits into the sum $\Delta_{2m,k} = \Delta_{2m,k}^+ \oplus \Delta_{2m,k}^-$, where $\Delta_{2m,k}^\pm$ are the eigenspaces of the endomorphism $\phi_{2m,k}(e_1 \cdot \dots \cdot e_{2m})$ to the eigenvalue $\pm i^{m+k}$. Let us denote by $u(\delta) \in \mathbb{C}^2$ the vector $u(\delta) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -\delta i \end{pmatrix}$, $\delta = \pm 1$, and let

$$u(\delta_1, \dots, \delta_m) = u(\delta_1) \otimes \dots \otimes u(\delta_m) \quad \delta_j = \pm 1.\tag{3}$$

Then $(u(\delta_1, \dots, \delta_m) \mid \prod_{j=1}^m \delta_j = \pm 1)$ is an orthonormal basis of $\Delta_{2m,k}^\pm$ with respect to the standard scalar product of $\mathbb{C}^{2^m}$.

4 General properties of twistor spinors

In this section we recall some properties of twistor spinors which we will need in the following calculations. For proofs see [BFGK91]. Let (M^n, g) be an oriented semi-Riemannian spin manifold of dimension $n \geq 3$. We denote by S the spinor bundle of (M^n, g) , by $\nabla^S : \Gamma(S) \rightarrow \Gamma(T^*M \otimes S)$ the spinor derivative given by the Levi-Civita connection of (M^n, g) and by $D : \Gamma(S) \rightarrow \Gamma(S)$ the Dirac operator on S . Let $p : TM \otimes S \rightarrow TM \otimes S$ be the projection onto the kernel of the Clifford multiplication μ . The *twistor operator*

$$\mathcal{D} : \Gamma(S) \xrightarrow{\nabla^S} \Gamma(T^*M \otimes S) \xrightarrow{g} \Gamma(TM \otimes S) \xrightarrow{p} \Gamma(\ker \mu)$$

is locally given by

$$\mathcal{D}\varphi = \sum_{j=1}^n \varepsilon_j s_j \otimes (\nabla_{s_j}^S \varphi + \frac{1}{n} s_j \cdot D\varphi),$$

where $(s_1, \dots, s_n)$ is a local orthonormal basis and $\varepsilon_j = g(s_j, s_j) = \pm 1$. A spinor field $\varphi \in \Gamma(S)$ is called *twistor spinor* if $\mathcal{D}\varphi = 0$.

Proposition 1 *For a spinor field $\varphi \in \Gamma(S)$ the following conditions are equivalent:*

1. φ is a twistor spinor.
2. φ satisfies the so-called twistor equation

$$\nabla_X^S \varphi + \frac{1}{n} X \cdot D\varphi = 0\tag{4}$$

for all vector fields X .

3. There exists a spinor field $\psi \in \Gamma(S)$ such that

$$\psi = g(X, X)X \cdot \nabla_X^S \varphi \quad (5)$$

for all vector fields X with $|g(X, X)| = 1$.

The dimension of the space $\mathcal{T}(M^n, g)$ of all twistor spinors is conformally invariant and bounded by

$$\dim \mathcal{T}(M^n, g) \leq 2 \cdot 2^{\lfloor \frac{n}{2} \rfloor}.$$

If (M^n, g) is simply connected and conformally flat, $\dim \mathcal{T}(M^n, g) = 2 \cdot 2^{\lfloor \frac{n}{2} \rfloor}$.

In particular, the twistor spinors on the semi-Euclidean space $\mathbb{R}^{n,k} := (\mathbb{R}^n, \langle \cdot, \cdot \rangle_{n,k})$ are the functions

$$\mathcal{T}(\mathbb{R}^{n,k}) = \left\{ \varphi \in C^\infty(\mathbb{R}^{n,k}, \Delta_{n,k}) \mid \varphi(x) = u + x \cdot v; \ u, v \in \Delta_{n,k} \right\}.$$

Let R be the scalar curvature and Ric the Ricci curvature of (M^n, g) . $K : TM \rightarrow TM$ denotes the $(1,1)$ -Schouten tensor of (M^n, g)

$$K(X) = \frac{1}{n-2} \left(\frac{R}{2(n-1)} X - \text{Ric}(X) \right).$$

Furthermore, let W be the $(4,0)$ -Weyl tensor of (M, g) and let us denote by the same symbol the corresponding $(2,2)$ -tensor field $W : \Lambda^2 M \rightarrow \Lambda^2 M$. Then we have

Proposition 2 *Let $\varphi \in \Gamma(S)$ be a twistor spinor. Then*

$$\nabla_X^S D\varphi = \frac{n}{2} K(X) \cdot \varphi, \quad (6)$$

$$W(\eta) \cdot \varphi = 0, \quad (7)$$

for all vector fields X and 2-forms η .

Finally, we recall two possibilities to obtain new manifolds with twistor spinors from a given one.

Let $(\tilde{M}^n, \tilde{g})$ be a simply connected parallelizable semi-Riemannian manifold and let $A \subset I(\tilde{M}, \tilde{g})$ denote a discrete subgroup of orientation preserving isometries of $(\tilde{M}, \tilde{g})$. We trivialize the spin structure of $(\tilde{M}, \tilde{g})$ with respect to a fixed global orthonormal basis field $\mathfrak{a} = (\mathfrak{a}_1, \dots, \mathfrak{a}_n)$. For $\gamma \in A$ we denote by $\Gamma(x) \in SO(n, k)$ the matrix of $d\gamma_x$ with respect to $\mathfrak{a}(x)$ and $\mathfrak{a}(\gamma(x))$. Then there are two lifts $\tilde{\Gamma}^\pm$ of Γ into $Spin(n, k)$

$$\begin{array}{ccc} & Spin(n, k) & \\ \tilde{\Gamma}^\pm \nearrow & \downarrow \lambda & \\ \tilde{M} & \xrightarrow{\Gamma} & SO(n, k) \end{array}$$

Let $\mathcal{E}(A)$ be the set of all left actions of A on $\tilde{M} \times Spin(n, k)$ such that

$$\varepsilon(\gamma)(x, a) = (\gamma(x), \varepsilon(\gamma, x) \cdot a) \quad \text{and} \quad \varepsilon(\gamma, x) = \tilde{\Gamma}(x)^\pm.$$

This set of left actions $\mathcal{E}(A)$ corresponds to the set of spin structures of the oriented semi-Riemannian manifold $M = \tilde{M}/A$. The spinor bundle on M corresponding to the spin structure $\varepsilon \in \mathcal{E}(A)$ is given by

$$S_\varepsilon = \left(\tilde{M} \times \Delta_{n,k} \right) /_\varepsilon$$

where $\varepsilon(\gamma)(x, v) = (\gamma(x), \varepsilon(\gamma, x) \cdot v)$ for all $\gamma \in A$, $(x, v) \in \tilde{M} \times \Delta_{n,k}$. Hence, the spinor fields on M corresponding to $\varepsilon \in \mathcal{E}(A)$ are given by the ε -invariant functions

$$\Gamma(S_\varepsilon) = C^\infty(\tilde{M}, \Delta_{n,k})^\varepsilon := \left\{ \varphi \in C^\infty(\tilde{M}, \Delta_{n,k}) \mid \varphi(\gamma(x)) = \varepsilon(\gamma, x) \cdot \varphi(x) \right\},$$

and for the twistor spinors on M with the spin structure ε we have

Proposition 3 *The twistor spinors on $M = \tilde{M}/A$ with respect to the spin structure $\varepsilon \in \mathcal{E}(A)$ are given by*

$$\mathcal{T}((M, g), \varepsilon) = \{ \varphi \in \mathcal{T}(\tilde{M}, \tilde{g}) \mid \varphi \text{ is } \varepsilon\text{-invariant} \}.$$

Let (M^{n+1}, g) be a semi-Riemannian spin manifold with spinor bundle S_M and let $F^n \subset M^{n+1}$ be a non-degenerate oriented hypersurface in M^{n+1} . We denote by $\eta : F \rightarrow TM$ the Gauss map of F , $\kappa(\eta) := g(\eta, \eta) = \pm 1$. It is well known, that the spinor bundle S_F of $(F, g|_F)$ with respect to the spin structure on F induced by the embedding, is isomorphic to $S_M|_F$ in case of even dimension n and to $S_M^\pm|_F$ in case of odd dimension n . Using this identification the Clifford multiplication and the spinor derivative are expressed by

$$\begin{aligned} X \cdot (\varphi|_F) &= (X \cdot \varphi)|_F \\ \nabla_X^{S_F}(\varphi|_F) &= \left(\nabla_X^{S_M^{(\pm)}} \varphi + \frac{1}{2} \kappa(\eta) \nabla_X^M \eta \cdot \eta \cdot \varphi \right) \Big|_F, \end{aligned}$$

where $\varphi \in \Gamma(S_M^{(\pm)})$, $X \in TF$ and $\varphi|_F$ always means the spinor field in $\Gamma(S_F)$ corresponding to φ with respect to the above mentioned isomorphism.

Proposition 4 *If $F^n \subset M^{n+1}$ is an umbilic hypersurface and $\varphi \in \Gamma(S_M^{(\pm)})$ is a twistor spinor on M , then $\varphi|_F \in \Gamma(S_F)$ is a twistor on F .*

Proof: Let $\lambda \in C^\infty(F)$ be the function satisfying $\nabla_X^M \eta = \lambda X$. Then

$$\nabla_X^{S_F}(\varphi|_F) = \left(\nabla_X^{S_M^{(\pm)}} \varphi + \frac{1}{2} \kappa(\eta) \lambda X \cdot \eta \cdot \varphi \right) \Big|_F.$$

If $\varphi \in \Gamma(S_M^{(\pm)})$ is a twistor spinor, from Proposition 1, (4), follows

$$\kappa(X) X \cdot \nabla_X^{S_M^{(\pm)}} \varphi = \frac{1}{n+1} D_M^{(\pm)} \varphi$$

for each $X \in TF$ with $\kappa(X) = g(X, X) = \pm 1$. Hence,

$$\kappa(X) X \cdot \nabla_X^{S_F}(\varphi|_F) = \left(\frac{1}{n+1} D_M^{(\pm)} \varphi - \frac{1}{2} \kappa(\eta) \lambda \eta \cdot \varphi \right) \Big|_F.$$

The right hand side is independent of $X \in TF$. Therefore, according to Proposition 1, (5), $\varphi|_F$ is a twistor spinor on F . $\square$

5 Twistor spinors on indecomposable, non-conformally flat Lorentzian symmetric spaces

Let us first consider the simply connected solvable Lorentzian symmetric space $M_{\underline{\lambda}}^n = (\mathbb{R}^n, g_{\underline{\lambda}})$, where

$$(g_{\underline{\lambda}})_{(s,t,x)} = 2 ds dt + \sum_{j=1}^{n-2} \lambda_j x_j^2 ds^2 + \sum_{j=1}^{n-2} dx_j^2,$$

and $\underline{\lambda} = (\lambda_1, \dots, \lambda_{n-2})$, $\lambda_i \in \mathbb{R} \setminus \{0\}$, $n \geq 3$. Let $\Lambda_0 := -\sum_{j=1}^{n-2} \lambda_j$. We fix the following global orthonormal basis on $M_{\underline{\lambda}}^n$:

$$\begin{aligned} \mathfrak{a}_{\overline{0}}(y) &:= \frac{\partial}{\partial s}(y) - \frac{1}{2} \left(\sum_{j=1}^{n-2} \lambda_j x_j^2 + 1 \right) \frac{\partial}{\partial t}(y) \\ \mathfrak{a}_0(y) &:= \frac{\partial}{\partial s}(y) - \frac{1}{2} \left(\sum_{j=1}^{n-2} \lambda_j x_j^2 - 1 \right) \frac{\partial}{\partial t}(y) \\ \mathfrak{a}_j(y) &:= \frac{\partial}{\partial x_j}(y) \quad j = 1, \dots, n-2, \end{aligned}$$

where $y = (s, t, x_1, \dots, x_{n-2}) \in M_{\underline{\lambda}}^n$. The vector field $V(y) := \frac{\partial}{\partial t}(y)$ is isotropic and parallel. The Ricci tensor of $M_{\underline{\lambda}}^n$ is given by

$$\text{Ric}(X) = \Lambda_0 \cdot g_{\underline{\lambda}}(X, V) V,$$

the scalar curvature R vanishes. Therefore, the Schouten tensor satisfies

$$K(X) = -\frac{1}{n-2} \Lambda_0 \cdot g_{\underline{\lambda}}(X, V) V. \quad (8)$$

For the Weyl tensor $W : \Lambda^2 M_{\underline{\lambda}} \rightarrow \Lambda^2 M_{\underline{\lambda}}$ one has

$$\begin{aligned} W(\mathfrak{a}_{\overline{0}} \wedge \mathfrak{a}_j) &= W(\mathfrak{a}_0 \wedge \mathfrak{a}_j) = (\lambda_j + \frac{1}{n-2} \Lambda_0) V \wedge \mathfrak{a}_j, \quad j = 1, \dots, n-2 \\ W(\mathfrak{a}_\alpha \wedge \mathfrak{a}_\beta) &= 0 \quad \text{for all other indices } \alpha, \beta, \end{aligned} \quad (9)$$

where $TM_{\underline{\lambda}}$ is identified with $T^*M_{\underline{\lambda}}$ using the metric $g_{\underline{\lambda}}$. In particular, $M_{\underline{\lambda}}$ is conformally flat if and only if $\underline{\lambda} = (\lambda, \dots, \lambda)$, $\lambda \in \mathbb{R} \setminus \{0\}$.

Since $M_{\underline{\lambda}}$ is simply connected, it has an uniquely determined spin structure. We trivialize this spin structure using the global orthonormal basis $(\mathfrak{a}_{\overline{0}}, \mathfrak{a}_0, \mathfrak{a}_1, \dots, \mathfrak{a}_{n-2})$ and identify the spinor fields with the smooth functions $C^\infty(M_{\underline{\lambda}}, \Delta_{n,1})$. The spinor derivative is defined by

$$\nabla_X^S \varphi = X(\varphi) + \frac{1}{2} \sum_{1 \leq k < l \leq n} \varepsilon_k \varepsilon_l g(\nabla_X^{LC} s_k, s_l) s_k \cdot s_l \cdot \varphi,$$

where $(s_1, \dots, s_n)$ is a local orthonormal basis and $\varepsilon_j = g(s_j, s_j) = \pm 1$. This gives for the spinor derivative on M_λ

$$\nabla_{\frac{\partial}{\partial t}}^S \varphi = \frac{\partial}{\partial t} \varphi \quad (10)$$

$$\nabla_{\frac{\partial}{\partial s}}^S \varphi = \frac{\partial}{\partial s} \varphi + \frac{1}{2} \sum_{j=1}^{n-2} \lambda_j x_j \mathbf{a}_j \cdot V \cdot \varphi \quad (11)$$

$$\nabla_{\frac{\partial}{\partial x_j}}^S \varphi = \frac{\partial}{\partial x_j} \varphi \quad (12)$$

$$\nabla_{\mathbf{a}_0}^S \varphi = \mathbf{a}_0(\varphi) + \frac{1}{2} \sum_{j=1}^{n-2} \lambda_j x_j \mathbf{a}_j \cdot V \cdot \varphi \quad (13)$$

$$\nabla_{\mathbf{a}_0}^S \varphi = \mathbf{a}_0(\varphi) + \frac{1}{2} \sum_{j=1}^{n-2} \lambda_j x_j \mathbf{a}_j \cdot V \cdot \varphi. \quad (14)$$

The vector space $\Delta_{n,1}$ is isomorphic to $\Delta_{n-2,0} \otimes \mathbb{C}^2$. Let us denote by Δ_V the subspace

$$\Delta_V := \Delta_{n-2,0} \otimes \mathbb{C}u(-1) \subset \Delta_{n,1}.$$

Using the formulas (1), (2) and (3) one obtains that a spinor field $\varphi \in C^\infty(M_\lambda, \Delta_{n,1})$ satisfies $V \cdot \varphi = 0$ if and only if the image of φ lies in Δ_V .

Proposition 5 *The space $\mathcal{P}(M_\lambda)$ of parallel spinors of M_λ is*

$$\mathcal{P}(M_\lambda) = \{ \varphi \in C^\infty(M_\lambda, \Delta_{n,1}) \mid \varphi = \text{constant} \in \Delta_V \}.$$

In particular, $\dim \mathcal{P}(M_\lambda) = \frac{1}{2} \cdot 2^{\lfloor \frac{n}{2} \rfloor}$.

Proof: From (10) - (12) it follows that $\varphi \in C^\infty(M_\lambda, \Delta_{n,1})$ is parallel if and only if φ depends only on s and

$$\frac{\partial \varphi}{\partial s} = -\frac{1}{2} \sum_{j=1}^{n-2} \lambda_j x_j \mathbf{a}_j \cdot V \cdot \varphi. \quad (15)$$

Therefore,

$$0 = \frac{\partial^2 \varphi}{\partial x_k \partial s} = -\frac{1}{2} \lambda_k \mathbf{a}_k \cdot V \cdot \varphi.$$

Since $\lambda_k \neq 0$ and $\mathbf{a}_k$ is space-like, this yields $V \cdot \varphi = 0$. Hence, because of (15), φ has to be constant. $\square$

Proposition 6 *Let M_λ be non-conformally flat. Then each twistor spinor on M_λ is parallel. In particular,*

$$\dim \mathcal{T}(M_\lambda) = \frac{1}{2} \cdot 2^{\lfloor \frac{n}{2} \rfloor}.$$

Proof: Let $\varphi \in C^\infty(M_{\underline{\lambda}}, \Delta_{n,1})$ be a twistor spinor. Then according to (7) of Proposition 2 $W(\eta) \cdot \varphi = 0$ for each 2-form η . Using (9) we obtain

$$\left(\lambda_j + \frac{1}{n-2} \Lambda_0\right) \mathfrak{a}_j \cdot V \cdot \varphi = 0 \quad j = 1, \dots, n-2.$$

Since $M_{\underline{\lambda}}$ is not conformally flat and $\mathfrak{a}_j$ is space-like, it follows

$$0 = V \cdot \varphi \tag{16}$$

Furthermore, we have

$$\nabla_X^S D\varphi \stackrel{(6)}{=} \frac{n}{2} K(X) \cdot \varphi.$$

Using (8) and (16) we obtain

$$\nabla_X^S D\varphi = -\frac{n}{2(n-2)} \Lambda_0 g_{\underline{\lambda}}(X, V) V \cdot \varphi = 0.$$

Hence, $D\varphi$ is parallel. From Proposition 5 follows, that $D\varphi$ is constant and $V \cdot D\varphi = 0$. Then the twistor equation and (10) yield

$$0 = \nabla_V^S \varphi + \frac{1}{n} V \cdot D\varphi = \frac{\partial}{\partial t}(\varphi).$$

Therefore, φ does not depend on t . Moreover, the twistor equation implies that

$$-\mathfrak{a}_{\overline{0}} \cdot \nabla_{\mathfrak{a}_{\overline{0}}}^S \varphi = \mathfrak{a}_0 \cdot \nabla_{\mathfrak{a}_0}^S \varphi = \frac{1}{n} D\varphi.$$

Then the formulas (13), (14) and (16) give

$$D\varphi = -n\mathfrak{a}_{\overline{0}} \cdot \left(\frac{\partial}{\partial s} - \frac{1}{2} \left(\sum_{j=1}^{n-2} \lambda_j x_j^2 + 1 \right) \frac{\partial}{\partial t} \right) (\varphi) = n\mathfrak{a}_0 \cdot \left(\frac{\partial}{\partial s} - \frac{1}{2} \left(\sum_{j=1}^{n-2} \lambda_j x_j^2 - 1 \right) \frac{\partial}{\partial t} \right) (\varphi).$$

Since φ does not depend on t we obtain

$$2D\varphi = n(\mathfrak{a}_0 - \mathfrak{a}_{\overline{0}}) \cdot \frac{\partial}{\partial s}(\varphi) = nV \cdot \frac{\partial}{\partial s}(\varphi) = n \frac{\partial}{\partial s}(V \cdot \varphi) \stackrel{(16)}{=} 0.$$

Therefore, φ is harmonic and the twistor equation implies that φ is parallel. $\square$

Now, let (M^n, g) be a non-conformally flat, non-simply connected, indecomposable Lorentzian symmetric space of dimension $n \geq 3$. Then, according to the Theorems 1,2 and 4, (M^n, g) is isometric to $M_{\underline{\lambda}}^n/A$, where A is a discrete subgroup of the centralizer $Z_{\underline{\lambda}} := Z_{I(M_{\underline{\lambda}})}(G(M_{\underline{\lambda}}))$.

1. case: There exist $i \in \{1, \dots, n-2\}$ such that $\lambda_i > 0$ or (i, j) such that $\frac{\lambda_i}{\lambda_j} \notin \mathbb{Q}^2$. Then $Z_{\underline{\lambda}} \simeq \mathbb{R} = \{\gamma_\alpha \mid \gamma_\alpha(s, t, x) = (s, t + \alpha, x), \alpha \in \mathbb{R}\}$.

Let $\gamma \in A$. With respect to the global basis $(\mathfrak{a}_{\overline{0}}, \mathfrak{a}_0, \mathfrak{a}_1, \dots, \mathfrak{a}_{n-2})$ the differential $d\gamma_y$ corresponds to the matrix $\Gamma(y) \equiv E \in SO(n, 1)$. Hence, $\tilde{\Gamma}^\pm(y) = \pm 1 \in Spin(n, 1)$. Therefore, we have 2 spin structures on $M = M_{\underline{\lambda}}/A$ corresponding to the homomorphisms $\text{Hom}(A; \mathbb{Z}_2)$. If $\varepsilon \in \text{Hom}(A, \mathbb{Z}_2)$ is not trivial, there are no non-trivial

ε -invariant constant spinor fields on $M_{\underline{\lambda}}$. From the Propositions 3,5 and 6 it follows that the twistor spinors on $M = M_{\underline{\lambda}}/A$ are given by

$$\mathcal{T}(M_{\underline{\lambda}}/A, \varepsilon) = \begin{cases} \{\varphi \in C^\infty(M, \Delta_V) \mid \varphi \text{ const}\} & \varepsilon \text{ trivial} \\ \{0\} & \varepsilon \text{ non-trivial.} \end{cases}$$

2. case: Let $\lambda_j = -k_i^2 < 0$ and $\frac{k_i}{k_j} \in \mathbb{Q}$ for all $i, j = 1, \dots, n-2$. Then

$$Z_{\underline{\lambda}} \simeq \left\{ \gamma_{\underline{m}, \alpha} \left| \begin{array}{l} \gamma_{\underline{m}, \alpha}(s, t, x) := (s + \beta, t + \alpha, (-1)^{m_1} x_1, \dots, (-1)^{m_{n-2}} x_{n-2}); \\ \text{where } \alpha \in \mathbb{R}, \underline{m} = (m_1, \dots, m_{n-2}) \in \mathbb{Z}^{n-2}, \\ \beta = \pi \cdot \frac{m_i}{k_i} \text{ } i = 1, \dots, n-2 \end{array} \right. \right\}$$

$$\simeq \mathbb{Z} \oplus \mathbb{R}.$$

A discrete subgroup $A_{\underline{m}, \alpha} \subset Z_{\underline{\lambda}}$ is generated by $\gamma_{\underline{m}, 0}$ and $\gamma_{0, \alpha}$. Let us suppose that $\sum_{i=1}^{n-2} m_i$ is even since otherwise $M_{\underline{\lambda}}/A_{\underline{m}, \alpha}$ is not orientable. $(d\gamma_{\underline{m}, \alpha})_y$ corresponds to the matrix

$$\Gamma(y) = \begin{pmatrix} 1 & & & & 0 \\ & 1 & & & \\ & & (-1)^{m_1} & & \\ & & & \ddots & \\ 0 & & & & (-1)^{m_{n-2}} \end{pmatrix}.$$

Hence $\tilde{\Gamma}^\pm(y) = \pm e_1^{m_1} \dots e_{n-2}^{m_{n-2}}$. Let $m_{i_1}, \dots, m_{i_s}$ be the odd elements in the tuple $\underline{m}$ ($s \in 2\mathbb{Z}$), and let us denote by $\omega_{\underline{m}} \in Spin(n-2, 0) \subset Spin(n, 1)$ the element

$$\omega_{\underline{m}} = e_{i_1} \dots e_{i_s}.$$

Then because of $\omega_{\underline{m}}^2 = (-1)^{\frac{s}{2}}$, $\omega_{\underline{m}}$ is an involution on $\Delta_{n-2, 0}$ if $s \equiv 0(4)$ and an almost complex structure if $s \equiv 2(4)$. The eigenspaces of $\omega_{\underline{m}}$ to the eigenvalues ± 1 and $\pm i$, respectively, have the same dimension (see formulas (1) and (2)).

The manifold $M_{\underline{\lambda}}/A_{\underline{0}, \alpha}$, $\alpha \neq 0$, has 2 spin structures and the twistor spinors are given as in case 1. $M_{\underline{\lambda}}/A_{\underline{m}, 0}$, $\underline{m} \neq \underline{0}$, has 2 spin structures, described by the homomorphisms $\varepsilon_\pm \in \text{Hom}(A_{\underline{m}, 0}, Spin(n-2, 0))$ given by $\varepsilon_\pm(\gamma_{\underline{m}, 0}) = \pm \omega_{\underline{m}}$. Then, according to the Propositions 3,5 and 6 the twistor spinors on $M = M_{\underline{\lambda}}/A_{\underline{m}, 0}$ are

$$\mathcal{T}(M_{\underline{\lambda}}/A_{\underline{m}, 0}, \varepsilon_\pm) = \left\{ \varphi \in C^\infty(M_{\underline{\lambda}}, \Delta_{n, 1}) \mid \begin{array}{l} \varphi(y) \equiv v \otimes u(-1); \\ \text{where } v \in \Delta_{n-2, 0} \text{ and } \omega_{\underline{m}} \cdot v = \pm v \end{array} \right\}.$$

Hence,

$$\dim \mathcal{T}(M_{\underline{\lambda}}/A_{\underline{m}, 0}, \varepsilon_\pm) = \begin{cases} 0 & \text{if } s \equiv 2(4) \\ \frac{1}{4} \cdot 2^{\lfloor \frac{n}{2} \rfloor} & \text{if } s \equiv 0(4). \end{cases}$$

The manifold $M_{\underline{\lambda}}/A_{\underline{m}, \alpha}$, $\underline{m} \neq \underline{0}, \alpha \neq 0$, has 4 spin structures corresponding to the homomorphisms $\varepsilon \in \text{Hom}(A_{\underline{m}, \alpha}, Spin(n, 1))$ given by $\varepsilon(\gamma_{\underline{m}, 0}) = \pm \omega_{\underline{m}}$, $\varepsilon(\gamma_{0, \alpha}) = \pm 1$.

Hence,

$$\mathcal{T}(M_{\underline{\lambda}}/A_{\underline{m},\alpha}, \varepsilon) = \begin{cases} \{0\} & \varepsilon(\gamma_{0,\alpha}) = -1 \\ \mathcal{T}(M_{\underline{\lambda}}/A_{\underline{m},0}, \varepsilon_{\pm}) & \begin{matrix} \varepsilon(\gamma_{0,\alpha}) = 1 \\ \varepsilon(\gamma_{\underline{m},0}) = \pm \omega_{\underline{m}} \end{matrix} \end{cases}$$

Summing up, we have

Proposition 7 *Let (M^n, g) be an indecomposable, non-conformally flat Lorentzian symmetric spin manifold of dimension $n \geq 3$. Then each twistor spinor is parallel and the dimension of the space of twistor spinors is*

$$\dim \mathcal{T}(M^n, g) = q \cdot 2^{\lfloor \frac{n}{2} \rfloor},$$

where $q = 0, \frac{1}{4}$ or $\frac{1}{2}$, depending on the fundamental group $\pi_1(M)$ and on the spin structure.

6 Twistor spinors on indecomposable conformally flat Lorentzian symmetric spaces of non-constant sectional curvature

According to the Theorems 1, 2 and 4 there are two isometry classes of indecomposable, conformally flat, simply connected Lorentzian symmetric spaces of dimension $n \geq 3$ and non-constant sectional curvature, namely

$$M_{\pm}^n := (\mathbb{R}^n, g_{\pm}),$$

where

$$(g_{\pm})_{(s,t,x)} = 2ds dt \pm ||x||^2 ds^2 + \sum_{j=1}^{n-2} dx_j^2.$$

Knowing that $\dim \mathcal{T}(M_{\pm}^n) = 2 \cdot 2^{\lfloor \frac{n}{2} \rfloor}$ we want to describe the twistor spinors explicitly. Let $w_1, w_2, w_3, w_4 \in \Delta_{n-2,0}$ and let us denote by $\varphi_{w_1, w_2, w_3, w_4} \in C^\infty(M_{\pm}^n, \Delta_{n,1})$ the following smooth functions

$$\begin{aligned} \varphi_{w_1, w_2, w_3, w_4}(s, t, x) &:= (\mp f'(s)w_3 - f(s)w_4 + x \cdot w_1) \otimes u(1) \\ &\quad + [-2w_1t + w_2 + x \cdot (f(s)w_3 + f'(s)w_4)] \otimes u(-1), \end{aligned}$$

where

$$f(s) = \begin{cases} \sinh(s) & \text{for } M_+^n \\ \sin(s) & \text{for } M_-^n. \end{cases}$$

Proposition 8 *The twistor spinors on $M_{\pm}^n$ are*

$$\mathcal{T}(M_{\pm}^n) = \{\varphi_{w_1, w_2, w_3, w_4} \mid w_1, w_2, w_3, w_4 \in \Delta_{n-2,0}\}.$$

Proof: We use the identification

$$\begin{aligned}\Delta_{n,1} &\simeq \Delta_{n-2,0} \otimes \Delta_{2,1} && \xrightarrow{\sim} \Delta_{n-2,0} \oplus \Delta_{n-2,0} \\ \varphi = \varphi_1 \otimes u(1) + \varphi_2 \otimes u(-1) &&& \longmapsto (\varphi_1, \varphi_2).\end{aligned}$$

Then according to (1) and (2) the Clifford multiplication corresponds to

$$\begin{aligned}X \cdot \varphi &= (-X \cdot \varphi_1, X \cdot \varphi_2) \quad \text{if } X \in \text{span}(\mathfrak{a}_1, \dots, \mathfrak{a}_{n-2}), \\ \mathfrak{a}_0 \cdot \varphi &= (-\varphi_2, -\varphi_1) \quad , \quad \mathfrak{a}_0 \cdot \varphi = (-\varphi_2, \varphi_1) \quad , \quad V \cdot \varphi = (0, 2\varphi_1).\end{aligned}$$

For the spinor derivative we obtain

$$\begin{aligned}\nabla_{\mathfrak{a}_0}^S \varphi &= (\mathfrak{a}_0(\varphi_1), \mathfrak{a}_0(\varphi_2) \pm x \cdot \varphi_1) \quad , \quad \nabla_{\mathfrak{a}_0}^S \varphi = (\mathfrak{a}_0(\varphi_1), \mathfrak{a}_0(\varphi_2) \pm x \cdot \varphi_1), \\ \nabla_{\mathfrak{a}_k}^S \varphi &= (\mathfrak{a}_k(\varphi_1), \mathfrak{a}_k(\varphi_2)) \quad , \quad k = 1, \dots, n-2.\end{aligned}$$

Let $\varphi = (\varphi_1, \varphi_2)$ be a twistor spinor on $M_{\pm}^n$. Then, according to Proposition 1, there exists a spinor field $\psi = (\psi_1, \psi_2)$ on $M_{\pm}^n$ such that

$$(\psi_1, \psi_2) = \mathfrak{a}_k \cdot \nabla_{\mathfrak{a}_k}^S \varphi = (-\mathfrak{a}_k \cdot \mathfrak{a}_k(\varphi_1), \mathfrak{a}_k \cdot \mathfrak{a}_k(\varphi_2)) \quad (17)$$

for each $k = 1, 2, \dots, n-2$. Therefore, $\varphi_1(s, t, \cdot)$ and $\varphi_2(s, t, \cdot)$ are twistor spinors on the Euclidean space $\mathbb{R}^{n-2}$. Hence,

$$\varphi_i(s, t, x) = u_i(s, t) + x \cdot v_i(s, t) \quad , \quad i = 1, 2,$$

where $u_i, v_i : \mathbb{R}^2 \rightarrow \Delta_{n-2,0}$. From (17) follows

$$\psi_1(s, t, x) = v_1(s, t) \quad \text{and} \quad \psi_2(s, t, x) = -v_2(s, t).$$

Furthermore, $\psi = (\psi_1, \psi_2)$ satisfies

$$(\psi_1, \psi_2) = -\mathfrak{a}_0 \cdot \nabla_{\mathfrak{a}_0}^S \varphi = \mathfrak{a}_0 \cdot \nabla_{\mathfrak{a}_0}^S \varphi.$$

Therefore,

$$v_1 = \left(\frac{\partial}{\partial s} - \frac{1}{2} (\pm \|x\|^2 + 1) \frac{\partial}{\partial t} \right) (u_2 + x \cdot v_2) \pm x \cdot (u_1 + x \cdot v_1) \quad (18)$$

$$v_1 = - \left(\frac{\partial}{\partial s} - \frac{1}{2} (\pm \|x\|^2 - 1) \frac{\partial}{\partial t} \right) (u_2 + x \cdot v_2) \mp x \cdot (u_1 + x \cdot v_1) \quad (19)$$

$$-v_2 = \left(\frac{\partial}{\partial s} - \frac{1}{2} (\pm \|x\|^2 + 1) \frac{\partial}{\partial t} \right) (u_1 + x \cdot v_1) \quad (20)$$

$$-v_2 = \left(\frac{\partial}{\partial s} - \frac{1}{2} (\pm \|x\|^2 - 1) \frac{\partial}{\partial t} \right) (u_1 + x \cdot v_1). \quad (21)$$

(18) + (19) gives $2v_1 = -\frac{\partial}{\partial t} u_2 - x \cdot \frac{\partial}{\partial t} v_2$ and after differentiation with respect to x_k $0 = -\mathfrak{a}_k \cdot \frac{\partial}{\partial t} v_2$. Hence,

$$\frac{\partial}{\partial t} v_2 = 0 \quad \text{and} \quad v_1 = -\frac{1}{2} \frac{\partial}{\partial t} u_2. \quad (22)$$

Using this, we obtain from (19) - (18)

$$0 = \mp x \cdot u_1 - \frac{\partial}{\partial s} u_2 - x \cdot \frac{\partial}{\partial s} v_2. \quad (23)$$

Differentiation and (22) show that

$$u_1 = \mp \frac{\partial}{\partial s} v_2 \quad \text{and} \quad \frac{\partial}{\partial t} u_1 = 0. \quad (24)$$

Inserting this in (23) and using (22) we obtain

$$\frac{\partial}{\partial s} u_2 = 0 \quad \text{and} \quad \frac{\partial}{\partial s} v_1 = 0. \quad (25)$$

Hence, $u_2 = u_2(t)$, $v_1 = v_1(t)$, $v_2 = v_2(s)$ and $u_1 = u_1(s)$. (21) - (20) shows

$$0 = x \cdot \frac{\partial}{\partial t} v_1 + \frac{\partial}{\partial t} u_1 = x \cdot v_1'(t).$$

Therefore, we have $v_1(t) \equiv w_1 \in \Delta_{n-2}$ and, because of (22), $u_2(t) = -2tw_1 + w_2$. (21) + (20) yields

$$-2v_2 = 2u_1'(s) \mp \|x\|^2 x \cdot v_1'(t) = 2u_1'(s),$$

so that, regarding (24), $v_2(s) = \pm v_2''(s)$. Therefore,

$$v_2(s) = f(s)w_3 + f'(s)w_4 \quad \text{and} \quad u_1(s) = \mp f'(s)w_3 - f(s)w_4,$$

where

$$f(s) = \begin{cases} \sinh(s) & \text{for } M_+^n \\ \sin(s) & \text{for } M_-^n. \end{cases}$$

Consequently, the twistor spinor φ is of the form $\varphi = \varphi_{w_1, w_2, w_3, w_4}$. $\square$

Now, let (M^n, g) be an indecomposable, conformally flat non-simply connected Lorentzian symmetric space of dimension $n \geq 3$ and non-constant sectional curvature. Then (M^n, g) is isometric to M_+^n/A or to M_-^n/A , where A is a discrete subgroup of

$$Z_+ := Z_{I(M_+)}(G(M_+)) = \{\varphi_\alpha \mid \varphi_\alpha(s, t, x) = (s, t + \alpha, x); \quad \alpha \in \mathbb{R}\}$$

in the first and of

$$Z_- := Z_{I(M_-)}(G(M_-)) = \left\{ \varphi_{m, \alpha} \mid \varphi_{m, \alpha}(s, t, x) = (s + m\pi, t + \alpha, (-1)^m x); \right. \\ \left. m \in \mathbb{Z}, \alpha \in \mathbb{R} \right\}$$

in the second case.

1. case: $M = M_+^n/A_\alpha$, $A_\alpha = \mathbb{Z}\varphi_\alpha$. Then there are 2 spin structures corresponding to $\varepsilon \in \text{Hom}(A_\alpha, \mathbb{Z}_2)$. The Propositions 3 and 8 show

$$\mathcal{T}(M, \varepsilon) = \begin{cases} \{\varphi_{0, w_2, w_3, w_4} \mid w_2, w_3, w_4 \in \Delta_{n-2, 0}\} & \varepsilon = 1 \\ \{0\} & \varepsilon \neq 1. \end{cases}$$

2. case: $M = M_-^n/A_{m, \alpha}$, $A_{m, \alpha} = \langle \varphi_{m, 0}, \varphi_{0, \alpha} \rangle$. If m is even and $\alpha \neq 0$, we have the same result as in case 1, since $f(s) = \sin(s)$ is $2\pi\mathbb{Z}$ -invariant. If m is odd, M

is orientable only if n is even. Then M^{2k} has 2 spin structures if $\alpha = 0$ and 4 spin structures if $\alpha \neq 0$. The Propositions 3 and 8 show

$$\mathcal{T}(M_-^{2k}/A_{m,0}, \varepsilon) = \begin{cases} \{0\} & n = 2k \equiv 0(4) \\ \left\{ \begin{array}{l} \varphi_{w_1, w_2, w_3, w_4} \mid \\ w_1, w_2 \in \Delta_{n-2,0}^\pm \\ w_3, w_4 \in \Delta_{n-2,0}^\mp \end{array} \right\} & \begin{array}{l} n = 2k \equiv 2(4) \\ \varepsilon(\varphi_{m,0}) = \pm e_1 \cdot \dots \cdot e_{n-2} \end{array} \end{cases}$$

$$\mathcal{T}(M_-^{2k}/A_{m,\alpha}, \varepsilon) = \begin{cases} \{0\} & \begin{array}{l} \varepsilon(\varphi_{0,\alpha}) = -1 \\ \text{or } n = 2k \equiv 0(4) \end{array} \\ \left\{ \begin{array}{l} \varphi_{0, w_2, w_3, w_4} \mid \\ w_2 \in \Delta_{n-2,0}^\pm \\ w_3, w_4 \in \Delta_{n-2,0}^\mp \end{array} \right\} & \begin{array}{l} n = 2k \equiv 2(4) \\ \varepsilon(\varphi_{0,\alpha}) = 1 \text{ and} \\ \varepsilon(\varphi_{m,0}) = \pm e_1 \cdot \dots \cdot e_{n-2} \end{array} \end{cases}$$

Summing up, we have in particular:

Proposition 9 *Let (M^n, g) be an indecomposable, conformally flat Lorentzian spin manifold (M^n, g) of non-constant sectional curvature and dimension $n \geq 3$. Then the dimension of the space of twistor spinors is*

$$\dim \mathcal{T}(M^n, g) = q \cdot 2^{\lfloor \frac{n}{2} \rfloor},$$

where $q = 0, \frac{3}{4}, 1, \frac{3}{2}$ or 2, depending on the fundamental group $\pi_1(M)$ and on the spin structure.

7 Twistor spinors on Lorentzian symmetric spaces of constant sectional curvature

Let $\psi_{u,v} \in C^\infty(\mathbb{R}^{n+1,k}, \Delta_{n+1,k})$ denote the twistor spinors on the pseudo-Euclidean space $\mathbb{R}^{n+1,k}$

$$\psi_{u,v}(x) := u + x \cdot v \quad , \quad u, v \in \Delta_{n+1,k}.$$

The pseudo-sphere $S_1^n(r) \subset \mathbb{R}^{n+1,1}$ and the pseudo-hyperbolic space $H_1^n(r)$ are umbilic hypersurfaces. Using the identification of the spinor bundle of the hypersurface with that of the external space (see section 4) we obtain from Proposition 4:

Proposition 10 *The twistor spinors on $S_1^n(r)$ and $H_1^n(r)$ with the induced spin structure are*

$$\mathcal{T}(S_1^n(r)) = \left\{ \psi_{u,v}|_{S_1^n(r)} \mid \begin{array}{ll} u, v \in \Delta_{n+1,1} & \text{if } n \equiv 0(2) \\ u \in \Delta_{n+1,1}^+, v \in \Delta_{n+1,1}^- & \text{if } n \equiv 1(2) \end{array} \right\}$$

$$\mathcal{T}(H_1^n(r)) = \left\{ \psi_{u,v}|_{H_1^n(r)} \mid \begin{array}{ll} u, v \in \Delta_{n+1,2} & \text{if } n \equiv 0(2) \\ u \in \Delta_{n+1,2}^+, v \in \Delta_{n+1,2}^- & \text{if } n \equiv 1(2) \end{array} \right\}.$$

The Lorentzian manifold $S_1^n(r)/\{\pm I\}$ is orientable if and only if n is odd, hence let n be odd. The volume form $\omega_{n+1,1} = e_1 \cdot \dots \cdot e_{n+1} \in Spin(n+1,1)$ satisfies $\omega_{n+1,1}^2 = (-1)^{\frac{n+1}{2}+1}$. Therefore, $S_1^n(r)/\{\pm I\}$ has no spin structure, if $n \equiv 3(4)$ and 2 spin structures, if $n \equiv 1(4)$. The spinor fields to these different spin structures can be identified with the invariant functions $C^\infty(S_1^n(r), \Delta_{n+1,1}^+)^{\varepsilon_\pm}$, where $\varepsilon_\pm$ is the $\mathbb{Z}_2$ -action given by

$$(\varepsilon_\pm(-1)\varphi)(x) = \pm\omega_{n+1,1} \cdot \varphi(-x) = \pm\varphi(-x).$$

From the Propositions 3 and 10 follows

$$\mathcal{T}(S_1^{4k+1}(r)/\{\pm I\}, \varepsilon) = \begin{cases} \{\psi_{u^+,0}|_{S_1^{4k+1}} \mid u^+ \in \Delta_{4k+2,1}^+\} & \text{if } \varepsilon = \varepsilon_+ \\ \{\psi_{0,v^-}|_{S_1^{4k+1}} \mid v^- \in \Delta_{4k+2,1}^-\} & \text{if } \varepsilon = \varepsilon_- \end{cases}$$

Now, let us consider a Lorentzian symmetric space M^n of constant negative sectional curvature. Then M^n is isometric to a Lorentzian covering of $H_1^n(r)/\{\pm I\}$. Let

$$\begin{aligned} \tilde{\pi} : \widetilde{H_1^n(r)} = \mathbb{R} \times \mathbb{R}^{n-1} &\longrightarrow H_1^n(r) \subset \mathbb{R}^{2,2} \times \mathbb{R}^{n-1} \\ (t, x) &\longmapsto (\sqrt{r^2 + \|x\|^2} \cos t, \sqrt{r^2 + \|x\|^2} \sin t, x) \end{aligned}$$

be the universal Lorentzian covering of $H_1^n(r)$. Let $\hat{Q}$ denote the reduction of the trivial spin structure Q of $\mathbb{R}^{n+1,2}$ to the subgroup $Spin(n,1)$ given by the Gaus map. Then $\hat{Q} := \tilde{\pi}^* \hat{Q}$ is the uniquely determined spin structure of $\widetilde{H_1^n(r)}$. The spinor fields of $\widetilde{H_1^n(r)}$ can be identified with the smooth functions $C^\infty(\widetilde{H_1^n(r)}, \Delta_{n+1,2}^{(\pm)})$, the twistor spinors are given by

$$\mathcal{T}(\widetilde{H_1^n(r)}) = \{\tilde{\psi}_{u,v} := \psi_{u,v}|_{H_1^n(r)} \circ \tilde{\pi} \mid \psi_{u,v}|_{H_1^n(r)} \in \mathcal{T}(H_1^n(r))\}.$$

Let

$$\begin{aligned} \pi_m : N_m^n &\longrightarrow H_1^n(r) \\ (\sqrt{\cdot} \cos t, \sqrt{\cdot} \sin t, x) &\longmapsto (\sqrt{\cdot} \cos(mt), \sqrt{\cdot} \sin(mt), x), \end{aligned}$$

$(\sqrt{\cdot} = \sqrt{r^2 + \|x\|^2})$, be the Lorentzian covering of $H_1^n(r)$ with respect to $m\mathbb{Z} \subset \pi_1(H_1^n(r)) = \mathbb{Z}$, $m = 1, 2, 3, \dots$. The manifold N_m^n has 2 spin structures. The corresponding spinor fields are given by the $\varepsilon_m^\pm$ -invariant functions $C^\infty(\widetilde{H_1^n(r)}, \Delta_{n+1,2}^{(\pm)})^{\varepsilon_m^\pm}$, where $\varepsilon_m^\pm$ is the $m\mathbb{Z}$ -action

$$(\varepsilon_m^\pm(mz)\varphi)(t, x) = (\pm 1)^z \varphi(t + 2\pi mz, x).$$

Therefore, the twistor spinors on N_m^n are

$$\mathcal{T}(N_m^n, \varepsilon) = \begin{cases} \{\psi_{u,v}|_{H_1^n(r)} \circ \pi_m \mid \psi_{u,v}|_{H_1^n(r)} \in \mathcal{T}(H_1^n(r))\} & \text{if } \varepsilon = \varepsilon_m^+ \\ \{0\} & \text{if } \varepsilon = \varepsilon_m^- \end{cases}$$

Finally, let us consider the manifolds $N_m^n/\{\pm I\}$. Since $N_m^n/\{\pm I\}$ is orientable if and only if n is odd, let n be odd. For the volume form $\omega_{n+1,2} = e_1 \cdot \dots \cdot e_{n+1} \in Spin(n+1,2)$ we have $\omega_{n+1,2}^2 = (-1)^{\frac{n+1}{2}+2}$. Therefore, there is no spin structure on $N_m^n/\{\pm I\}$ if

$n \equiv 1(4)$ and there are 4 spin structures in case $n \equiv 3(4)$. The spinor fields are given by the functions $C^\infty(\widetilde{H_1^n(r)}, \Delta_{n+1,2}^+(\varepsilon_m^\pm, \delta^\pm))$, invariant under the $m\mathbb{Z}$ -action $\varepsilon_m^\pm$ and the $\mathbb{Z}_2$ -action $\delta^\pm$ given by

$$\begin{aligned} (\delta^\pm(-1)\varphi)(t, x) &= \pm\omega_{n+1,2} \cdot \varphi(t + m\pi, -x) \\ &= \pm\varphi(t + m\pi, -x). \end{aligned}$$

Then, the twistor spinors are

$$\mathcal{T}\left(N_m^{4k+3}/\{\pm I\}, \varepsilon\right) = \begin{cases} \{0\} & \varepsilon = (\varepsilon_m^-, \delta^\pm) \text{ or} \\ & \varepsilon = (\varepsilon_m^+, \delta^-), m \equiv 0(2) \\ \{\tilde{\psi}_{u^+,0} \mid u^+ \in \Delta_{4k+4,2}^+\} & \varepsilon = (\varepsilon_m^+, \delta^+) \\ \{\tilde{\psi}_{0,v^-} \mid v^- \in \Delta_{4k+4,2}^+\} & \varepsilon = (\varepsilon_m^+, \delta^-), m \equiv 1(2). \end{cases}$$

Summing up, we have in particular

Proposition 11 *Let (M^n, g) be a Lorentzian symmetric spin manifold of constant sectional curvature $k \neq 0$ and dimension $n \geq 3$, then the dimension of the space of twistor spinors is*

$$\dim \mathcal{T}(M^n, g) = q \cdot 2^{\lfloor \frac{n}{2} \rfloor},$$

where $q = 0, 1$, or 2 depending on $\pi_1(M)$ and on the spin structure.

References

- [ACDS97] D.V. Alekseevsky, V. Cortes, C. Devchand, and U. Semmelmann. Killing spinors are Killing vector fields in Riemannian supergeometry. MPI-preprint 97-29, 1997.
- [Bau97] H. Baum. Strictly pseudoconvex spin manifolds, Fefferman spaces and Lorentzian twistor spinors. SFB 288-Preprint No. 250, 1997.
- [BFGK91] H. Baum, T. Friedrich, R. Grunewald, and I. Kath. *Twistors and Killing Spinors on Riemannian Manifolds*, volume 124 of *Teubner-Texte zur Mathematik*. Teubner-Verlag, Stuttgart/Leipzig, 1991.
- [CK78] M. Cahen and Y. Kerbrat. Champs de vecteurs conformes et transformations conformes des espaces Lorentzian symetriques. *J.Math. pures et appl.*, 57:99–132, 1978.
- [CLP⁺90] M. Cahen, J. Leroy, M. Parker, F. Tricerri, and L. Vanhecke. Lorentzian manifolds modelled on a Lorentz symmetric space. *J. Geom. Phys.*, 7(4):571–581, 1990.
- [CP80] M. Cahen and M. Parker. Pseudo-Riemannian symmetric spaces. *Mem. AMS*, 24(229):1–108, 1980.

- [CW70] M. Cahen and N. Wallach. Lorentzian symmetric spaces. *Bull. AMS*, 76(3):585–591, 1970.
- [Fri97] Th. Friedrich. *Dirac-Operatoren in der Riemannschen Geometrie*. Vieweg-Verlag, 1997.
- [Lew91] J. Lewandowski. Twistor equation in a curved spacetime. *Class. Quant. Grav.*, 8:11–17, 1991.
- [NW84] P. Nieuwenhuizen and N.P. Warner. Integrability conditions for Killing spinors. *Comm. Math. Phys.*, 93:277–284, 1984.
- [Pen67] R. Penrose. Twistor algebra. *J. Math. Phys.*, 8:345–366, 1967.
- [PR86] R. Penrose and W. Rindler. *Spinors and Space-time II*. Cambr. Univ. Press, 1986.
- [Wol84] J.A. Wolf. *Spaces of constant curvature*. Publish or Perish, Inc., 1984.

Helga Baum
 Institut für Reine Mathematik
 Humboldt-Universität zu Berlin
 Sitz: Ziegelstr. 13a
 10099 Berlin , Germany
 email: baum@mathematik.hu-berlin.de